

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/355792529>

Review On Biology, Distribution And Conservation Challenges For Horseshoe Crabs In India

Article in *Journal of Sustainability Science and Management* · October 2021

DOI: 10.46754/jssm.2021.10.023

CITATIONS

4

READS

903

10 authors, including:



Ayaskanta Pramanik

National Centre For Cell Science, Pune

2 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



Mr. Apratim Sai Rajesh

Fakir Mohan University

21 PUBLICATIONS 44 CITATIONS

[SEE PROFILE](#)



Sanatan Tudu

Fakir Mohan University

9 PUBLICATIONS 24 CITATIONS

[SEE PROFILE](#)



Melissa B. Martin

Australian Maritime College

25 PUBLICATIONS 240 CITATIONS

[SEE PROFILE](#)

REVIEW ON BIOLOGY, DISTRIBUTION AND CONSERVATION CHALLENGES FOR HORSESHOE CRABS IN INDIA

AYASKANTA PRAMANIK¹, APRATIM SAI RAJESH^{1,2}, SANATAN TUDU^{1,2}, MELISSA BEATA MARTIN⁶, SALWA SHAHIMI⁶, BRYAN RAVEEN NELSON^{2,3}, TANMAY SARKAR⁴, BEHARA SATYANARAYANA⁷, SIDDHARTHA PATI^{1,2*} AND BISNU PRASAD DASH^{1,2*}

¹Department of Biosciences and Biotechnology, Fakir Mohan University, Odisha-756089, India. ²SLAN Institute-Association for Biodiversity Conservation and Research (ABC), Odisha, 756001, India. ³Institute of Tropical Biodiversity and Sustainable Development, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia. ⁴Department of Food Technology and Bio-chemical Engineering, Jadavpur University, Jadavpur, Kolkata-700032, India. ⁵Malda Polytechnic, West Bengal State Council of Technical Education, Government of West Bengal, Malda-732102, India.

⁵Faculty of Science and Marine Environment, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

⁶Institute of Oceanography and Environment, Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

*Corresponding author: patisiddhartha@gmail.com/bisnubsbtjmu@gmail.com

Submitted final draft: 4 January 2021

Accepted: 26 January 2021

<http://doi.org/10.46754/jssm.2021.10.023>

Abstract: Although horseshoe crabs have ecological and biomedical value, their restricted distribution in the northeast coast (Odisha, West Bengal) to a limited number of researchers have made them an understudied arthropod in India. With an objective that uses 'desktop review' for horseshoe crab ecology, biology and their life cycle in India, it is learnt that actions to conserve *T. gigas* and *C. rotundicauda* are still in implementing stages. In Asia, *Tachypleus gigas* and *Carcinoscorpius rotundicauda* are present in sandy and muddy intertidal zones, which are termed nursery and feeding grounds in West Bengal, India. Broad tolerance to temperature (19-41°C) and salinity (2-36‰) variations indicate a non-paused horseshoe crab presence and reproduction for West Bengal. Horseshoe crabs are not harvested for human consumption in India but, industrial demands for embryonic perivitelline (growth enhancer and stimulus for stem cell differentiation) and novel therapeutic applications (from anti-microbial and anti-carcinogenic proteins) are present in small scales. Although horseshoe crabs are safeguarded by the Wildlife Protection Act, 1972, the definitions are limited to *T. gigas*. Therefore, the output of this study contributes data for IUCN Red List assessments because fisheries records have revealed novel horseshoe crab distribution sites, incidences of by-catch, poor by-catch management and also coastal interventions that risk horseshoe crab nursery grounds in India into a deletion.

Keywords: Arthropod, ecology, fisheries, habitat, industry, strategy.

Introduction

The *Tachypleus gigas* (Müller) and *Carcinoscorpius rotundicauda* (Latreille) are restricted to northeast India (Odisha, West Bengal) where they co-exist in sandy and muddy intertidal areas (Annandale, 1909; Rao & Rao, 1972; Lazarus *et al.*, 1990; Tripathy *et al.*, 2013; Yennawar, 2015; Pati *et al.*, 2015). Yet, intertidal zones should not be assumed as horseshoe crab habitat but rather, an indication of natal-homing into nursery grounds for their routine spawning (Nelson *et al.*, 2015; 2016a; b; Fairuz-Foziet *et al.*, 2018; John *et al.*, 2018; Nelson *et al.*, 2019; Zauki *et al.*, 2019a; b). In fact, horseshoe crabs were witnessed to co-exist in mangrove coasts of

Sundarban, Hukitola and Dhamra because sand, clay and silt in the ratio of 7:2:1 are preferred by both species for spawning (Chatterji *et al.*, 1992a; Chatterji *et al.*, 1996a). Research developments has updated existing knowledge on spatial-temporal patterns with transitional weather, population size, spawning cycles, habitat mapping and spawning migrations (Annandale, 1909; Seawell, 1912; Rao & Rao, 1972; Debnath *et al.*, 1989; Lazarus *et al.*, 1990; Chatterji *et al.*, 1992b; Chatterji *et al.*, 1996b; Chatterji, 1999; Chatterji.&Shaharom, 2009; Tripathy *et al.*, 2013; Yennawar, 2015; Pati *et al.*, 2015), embryonic development of *T. gigas* and *C. rotundicauda* (Roonwall, 1944; Debnath,




Figure 1: Horseshoe crab spawning and nursery grounds in West Bengal (India). The horseshoe crabs are abbreviated as TG = *Tachypleus gigas* and CR = *Carcinoscorpius rotundicauda*

1991; Chatterji *et al.*, 1992b; Chatterji *et al.*, 1996a; Khan, 2003; Chatterji *et al.*, 2004a), dietary and feeding habits (Debnath *et al.*, 1989; Chatterji *et al.*, 1992c) and exploitation of horseshoe crabs (Debnath & Choudhury, 1988; Pati & Dash, 2016).

After several man-made activities, like the excavation of sand, construction of jetties on seashore areas, the shore geomorphology (<1 m deep) was composed of different proportions of fine sand, and abrupt sediment nomenclature changes were decreasing the spawning emergence of horseshoe crabs in West Bengal over the last two decades (Chatterji *et al.*, 2004b; Chatterji & Shaharom, 2009; Behera *et al.*, 2015; Pati *et al.*, 2017). Although the

circulation of fine sand in Sundarban, Hukitola and Dhamra coasts is under control during the ‘no-fishing season’ under Indian Fisheries Act 1897 regulations, shore rehabilitation is a present challenge because local communities are regularly using estuaries for capture fisheries and recreational activities (Pati *et al.*, 2017). With horseshoe crabs included in Schedule IV of the Wildlife Protection Act 1972, these arthropods are treated as by-catch in India (Chatterji, 1999; Behera *et al.*, 2015). Therefore, researchers excluded capture fisheries from their attention and focused on horseshoe crab population assessments by comparing their spawning emergence during different seasons (John *et al.*, 2018). A new finding emerged, where *T. gigas*

preferred summer months (March-May, salinity 25-33 ‰) whereas *C. rotundicauda* gravitated towards the monsoon season (June-September, salinity 0-16 ‰) for peak spawning activity (Behera et al., 2015; Techera, 2020; Table 1).

Industrial activities have expanded the conversion of marine resources into products like health tonics and handicrafts, particularly in West Bengal (Northeast India) where traditional practices remain prevalent (Mondayil et al., 2006; Mondal & Bandyopadhyay, 2014; Pati et al., 2020a). For instance, horseshoe crab carapace oil is used as an aphrodisiac (Mishra, 2009a, b), whereas the powdered carapace is effective for lacerations, burn wounds, skin ailments and for food preservation (Kumar et al., 2015; Pati et al., 2018; Ibrahim et al., 2019; Pati et al., 2020b). Meanwhile, the perivitelline fluid that buffers *T. gigas* embryo from the external environment was fractionated to derive the presence of proteins (hemagglutinin, hemocyanin, 27 kDa lectin and L6 subunit), peptides and buffer salts (Mirshahi et al., 2011). Considering that perivitelline fluid function is similar to amniotic fluid, it can trigger the development of epithelial cells as well as clot blood (c.f. Oppenheim et al., 1974; Weaver et al., 1988). Modern applications include the use of *T. gigas* embryonic perivitelline fluid to enhance cell growth, ionic regulation (for biological filtering) and treat cardiovascular diseases, while its hemolymph is used as a tumor marker, indicate endotoxin presence and as an antibiotic (Srimal et al., 1985; Ghaskadbi et al., 2008; Mirshahi et al., 2011; Pati et al., 2015; Chinnari et al., 2015; Nelson et al., 2020; Nong et al., 2020).

Unfortunately, nursery and feeding grounds in India have been degraded, whereby the poor environmental health impacts the growth and development of larvae (Pati et al., 2021). Therefore, researchers were collecting *T. gigas* with weight (0-130 g), size (0-20 % w/w), density ($\pm 3 \text{ g cm}^{-1}$) and haemolymph viability (70-88 %) in negative trends from the same sources (Chatterji et al., 1988; 1992; 1994; Joshi et al., 2002; Vijayakumar et al., 2000; Sahu & Dey, 2013; Pati et al., 2018). With an

objective to compile information on horseshoe crab ecology, biology and life cycle, the findings were used to address conservation challenges for *T. gigas* and *C. rotundicauda* in India. Since *L. polyphemus* and *T. tridentatus* are listed as “Vulnerable” and “Endangered” while both, *T. gigas* and *C. rotundicauda* remain as “Data Deficient”, information on novel horseshoe crab distribution adds to existing data for the Red List (IUCN) assessment of these arthropods.

Methodology

Through a ‘desktop review’, words like horseshoe crabs, research, India, Bengal, northeast, West, conservation, threats, biology, spawning behaviour, *Tachypleus gigas* and *Carcinoscorpius rotundicauda* were used (independent or in combination) to search for literature in Google Scholar, Scopus, Web of Science, PubMed and NCBI. The library construction adopts the practice of John et al. (2018) where the inclusion criteria comprise published materials in journals, reports, book chapters, abstracts, symposium transcripts and theses between January 1900 and July 2020, and having touchpoints on horseshoe crabs from India. Yielding more than 100 published materials, an exclusion criterion focusing on biology, distribution and conservation challenges further limited the final library to only 56 information sources, which are available in this review.

Results and Discussions

Horseshoe Crab Ecology

Horseshoe crabs were studied for their distribution, biology, ecology, spawning behaviour through the 56 published materials in this review. In fact, the first publication was released by the National Museum about a strange crab called *L. moluccanns* (c.a. *T. gigas*) that occupies a 700 km continental shelf stretch situated between Bhadrak and Kendrapada (Odisha, East India) where depths are 20 fathoms (30-40 m). This arthropod emerges into shallow water in pairs during rising tides and the larger crab carries pale-yellow eggs




Figure 2: Horseshoe crab spawning and nursery grounds in Odisha (India). The horseshoe crabs are abbreviated as TG = *Tachypleus gigas* and CR = *Carcinoscorpius rotundicauda*

in the ventral section of its head (Annandale, 1909). Comparatively, the *C. rotundicauda* is described as a benthic crab that lingers in shallow and moderately deep (4-10 m) freshwater habitats some 60 miles upstream the Hoogly River (Seawall, 1912; Figs. 1-2). The taxonomic description of *L. moluccanns* was revised to *T. gigas* in 1944 and it was locally called *raj kakraor* ‘king crab’ for its larger appearance than decapod crustaceans (Roonwal, 1944). Later research discovered that *T. gigas* spawning activity in northeast Bengal coincides with one circa tidal lunar clock, the full moon ebb tide (Rao & Rao, 1972; Sekiguchi *et al.*, 1976). Then, hydrographic definitions like wave characteristics, seabed slope, and near-shore gradient were integrated into lunar ebb tide events as a response to enhance understanding on *T. gigas* spawning and larval recruitment (Chatterji *et al.*, 1996b).

A decline in adult *T. gigas* retrieval over a series of population assessment studies

motivated the inclusion of horseshoe crabs (specifically *T. gigas*) into schedule IV of the Wildlife Protection Act 1972 (Chatterji, 1999; Behera *et al.*, 2015). Separately, laboratory experiments on adult *T. gigas* revealed that low salinity (2-16 ‰) maintained their body weights but, reduced their haemocyte:haemolymph composition (viability) by 8 % (male) and 10 % (female) (Chatterji *et al.*, 1992b). Also, an *ex-situ* assay involving *T. gigas* trilobite larvae showed that 10 ‰ and 40 ‰ salinity increased the rate of embryogenesis by 7 days if compared to ambient (20-33‰) gradients (Chatterji *et al.*, 2004a). Together, these experiments related ecology conditions with the quality of crabs, particularly during the sourcing of horseshoe crabs for haemolymph and chitosan extraction (Pati *et al.*, 2018). In addition, research collaboration between India and Malaysia produced novel understanding for relationships between riverine mangrove vegetation, bedload and fluvial sediment transport, the relationships between sediment compaction and nest depth,

Table 1: The spawning ground conditions in Odisha (Saha, 2010)

Species	Area	Salinity (%)	pH	Nest (cm)		Egg (nos.)	Egg Size (mm)	Larva
				Width	Depth			
<i>T. gigas</i>	Sandy	11-36	7.29-8.35	12-30	10-30	60-720	3.7	Trilobites swimming to the sea with the low tide
<i>C. rotundicauda</i>	Mangrove & muddy	2-33	6.90-7.55	-	3-7	80-200	2.3	Juveniles found in the mangroves mud flat

the importance of osmotic and temperature shocks and also methods to identify *C. Rotundicauda* nests (Nelson *et al.*, 2015; Pati *et al.*, 2017; Fairuz-Fozi *et al.*, 2018; Nelson *et al.*, 2019; 2020).

Horseshoe Crab Biology

The anatomy of horseshoe crabs was first described using *L. polyphemus* (Heinbecker, 1932; Shuster Jr., 1948; Dumont *et al.*, 1965). Later revisions used *T. gigas* for mid-gut gland labels and also classification of its J-shaped digestive tract into fore-gut (mouth, oesophagus and proventriculus), mid-gut (comprising stomach and intestine with hepatic (yellow) connective tissue extending to the opisthosoma, whereas the diverticulum (branching hepatic ducts) almost occupies the entire prosoma) and hind-gut (rectum and anus) (Debnath *et al.*, 1989). Separately, *T. gigas* is recognized by its deep green shell with depressed lateral spines, six opisthosoma spines (equally long = male; 3 long and 3 short = female), movable spur on the 4th segment of the sixth prosomatic appendage and a triangular-shaped telson, whereas the *C. rotundicauda* has a deep brown-green shell, six prosomal spines of equal length for both genders, spurs are absent on the 4th segment of the sixth prosomatic appendage, its genital operculum extends distally to the tip of gill lamellae and has a smooth-rounded telson (Debnath, 1991).

With the discovery of histamine as a neurotransmitter in *L. polyphemus* (Battelle *et al.*, 1991), the *T. gigas* brain (collar around the oesophagus) was examined and classified into anterior (suprapharyngeal ganglion) and posterior (suprapharyngeal tritocerebrum ganglion) nerve cords where olfactory nerves connect to lateral eyes, and cheliceral nerves extend communication to its appendage (Debnath, 1991). Neurological and anatomical descriptions of *T. gigas* were later used to completely describe the body plan of *L. polyphemus* in Walls *et al.* (2002). The *T. gigas* feeds on decayed organic matter, detritus, scaphopods, gammarid amphipods, foraminifera, vascular plant, molluscs, polychaetes and crustaceans, all of which relate to its benthic lifestyle (Debnath *et al.*, 1989; Table 2). The presence of acid protease (gizzard>oesophagus), alkali protease (intestine>gizzard), esterase and cellulose (oesophagus), invertase and amylase (digestive gland) suggests that *T. gigas* is a generalist that can digest plant and animal materials (Debnath *et al.*, 1989; Chatterji *et al.*, 1992c). This arthropod does not have fixed diets and it would feed more during the winter months (December-January) (Nair *et al.*, 1990).

Relative growth relationships between length-weight and length-length were used to indicate *T. gigas* health (or wellbeing) and to identify mature size-classes in their native habitat (Chatterji *et al.*, 1988; 1992; 1994). Yet, merging horseshoe crab samples of different size

Table 2: Food sources discovered from the *Tachypleus gigas* gut (Debnath, 1989)

Bivalves	Gastropod	Annelids	Arthropods	Plant	Miscellaneous	Teleosts
<i>Anadra</i> sp.	<i>Cerithedia</i> sp.	<i>Gattyana</i> sp.	Mysid shrimps	<i>Casuarina</i> needles	<i>Ammonia beccarii</i>	Goboid fish
<i>Dosinia</i> sp.	<i>Littorina</i> sp.	<i>Phyllodoce</i> sp.	Gammarid amphipods	Plant fruits	<i>Asteroroatalia dentate</i>	
<i>Placenta</i> sp.	<i>Assimenea</i> sp.	<i>Nereis</i> sp.	Copepodas (<i>Microstella</i> sp.)	Algal bodies	<i>A. multispinosa</i>	
<i>Macoma</i> sp.	<i>Nerita</i> sp.	<i>Perinereis</i> Sp.	Decapods (<i>Crangon</i> sp.)		<i>A. trispinosa</i>	
<i>Solen</i> sp.	<i>Cassibula</i> sp.	<i>Glyceria</i> sp.	Cirripeds		<i>Bolivina</i> sp.	
<i>Neosolen</i> sp.	<i>Cymia</i> sp.	<i>Sabellaria</i> sp.	<i>Chrionomus</i> sp.		<i>Elphidium</i> sp.	
<i>Teredo</i> sp.		<i>Polydora</i> sp.	<i>Atylotus agrestis</i>		<i>Ouinquoloculina</i> sp.	

classes (male and female) and maturity (juvenile, sub-adult and adult) was a common practice and therefore, it implicated the accuracy results for weight ($\pm 24\%$) and size ($\pm 21\%$) analysis (Sahu & Dey, 2013). The male *L. polyphemus* is smaller than the female because it lacks the 17th moult stage (Levin, 2003). Therefore, male horseshoe crabs become sexually mature earlier than the female crabs, although they weigh much less than their counterparts (Chatterji *et al.*, 1988; Graham *et al.*, 2009). With weight influenced by eggs (for gravid female crabs) and feeding, morphometry studies indicated that prosoma of *T. gigas* develops symmetrically ($p = 0.87-0.95$) with its total length (Vijayakumar *et al.*, 2000). Altogether, the *T. gigas* relative growth assessments were later revised by sub-clustering samples into different gender, size class and habitat groups (Vijayakumar *et al.*, 2000; Khan, 2003; Sahu & Dey, 2013; Chatterji & Pati, 2014; Panda & Naik, 2017). Novelty by studies were recognized and adapted with *T. tridentatus*, *L. polyphemus* and *T. gigas* morphology assessments in Asia (Chiu & Morton, 2003; Srijaya *et al.*, 2010; Smith & Brockmann, 2014; Kumar *et al.*, 2015; Mohamad *et al.*, 2016; Jawahir *et al.*, 2017; Razak & Kassim, 2018; Putri *et al.*, 2019; Syuhaida *et al.*, 2019).

Life Cycle

Researchers learnt that horseshoe crab size varies by gender, but confusion arises when same size and different gender crabs from both species were presented together. It was resolved when researchers compared the hook-like structure of 2nd and 3rd appendages in which, only male crabs possess a degenerate propodiate while the dactylopodiate appears similar for all appendages (Rao & Rao, 1972). Species-wise, male *T. gigas* possesses an underdeveloped propodiate and its 2nd and 3rd opisthosoma marginal spines are slightly longer than other spines, whereas the *C. rotundicauda* has a well-developed propodiate and its opisthosoma marginal spines are arranged in reducing order for their length (Debnath, 1991). The marked morphological differences between male and female could also indicate their maturity, where matured crabs appear in dark tans, mating scars and broken spines are indications of sexually active females, while male crabs have bulged lateral prosoma flesh (Debnath & Choudhury, 1988). Though widely accepted that horseshoe crabs will become encrusted by fouling communities if the crab remains dormant for long periods (John *et al.*, 2018), it was previously assumed that epifauna (fouling

Table 3: Epifauna that were fouling the *Tachypleus gigas* carapace in India (Debnath, 1989)

Species Name	Class/Order	<i>T. gigas</i>	<i>C. rotundicauda</i>
<i>Metridium schillerianum</i>	Anthozoa	+	+
<i>Gattyana deludens</i>	Polychaeta	+	
<i>Balanus darwini</i>	Hexanauplia	+	+
<i>Chthamalus stellatus</i>	Hexanauplia		+
<i>Cheiriphotis megacheles</i>	Amphipoda	+	
<i>Cleantis natalensis</i>	Isopoda	+	
<i>Ostrea sp.</i>	Bivalvia		+

community) encrusting is a parasite infection (Roonwal, 1944; Rao & Rao, 1972). Later, *ex-situ* experiments showed that epifauna-infected crabs were less active and inspection of genital pores after epifauna removal led researchers to conclude that only infertile mature crabs become inhabited by fouling communities (Debnath & Choudhury, 1991; Patil & Anil, 2000; Table 3).

Meanwhile, the absence and presence of nests, closely spaced nests and >1 egg clutch per nest indicated the possibility of multiple spawning incidences per amplexus, fecundity to relative size and also spawning migration (Chatterji & Parulekar, 1992; Chatterji *et al.*, 1992; 1996). On the other hand, carapace-like impressions, nest diameter (14-30 cm), nest depth (2-3 inches) and attaining 20-100 eggs per clutch, were indications that photoperiod, tidal and lunar cycles are influencing the spawning behaviour of *T. gigas* during the summer months (March-May) and *C. Rotundicauda* during the

monsoon season (June-September) in northeast India (Debnath & Choudhury, 1988; Debnath & Choudhury, 1992; Chatterji *et al.*, 1992a, b; Mishra *et al.*, 1993; Sahu & Dey, 2013; Chatterji *et al.*, 1996b; Figs. 1-2). Considering that transitional weather shifts shore geomorphology (0.182-0.203 mm) and sediment temperature (31.8-34.6°C), disparities in spawning yields (eggs and nests) indicate the possibility of spawning migration for *T. gigas* (Chatterji *et al.*, 1996b; Chatterji, 1999), whereby *T. gigas* increases its home range beyond the natal shore boundary (Tripathy *et al.*, 2013).

The eggs of *T. gigas* are green, spherical, protected by a thin elastic euticulablastodermica, encapsulated by a leathery chorion and have nine embryogenesis stages before hatching of the trilobite larvae (Roonwal, 1944). Later, a total of four embryonic moults were discovered for *L. polyphemus* (Bennett *et al.*, 1972; Bannon & Brown, 1980) and *T. tridentatus*, including the hatching that commences after 25-32 days of embryogenesis (Yamamichi & Sekiguchi, 1974; Yamamichi *et al.*, 1983). Annotations on *T. gigas* embryogenesis by Chatterji *et al.* (1996a) reveals that newly fertilized eggs (size 3.57 mm and weight 24.17 mg; Stage A), rupturing of chorion that enlarges the egg to 3.86 mm (Stage C; Moults 1), separation of leathery chorion after swelling of euticulablastodermica by perivitelline fluid build-up (Stage D; Moults 2), embryonic cleavage for vasculogenesis (Stage E; Moults 3), embryo weighs 138.02 mg and has movable legs (Stage F) and hatching of trilobite larvae that measures 6.99 mm and weighs



Figure 3: Liberation of horseshoe crab (*Tachypleus gigas*) oil by boiling the carapace

156.22 mm (Stage I; Moults 4). This annotation considers all nine development stages described by Roonwall (1944) and adopts the embryonic moults described by Yamamichi *et al.* (1983) to produce complete understanding on *T. gigas* embryonic life stages.

Conservation Challenges

The dorsal shell (exoskeleton) of horseshoe crabs is armoured by a chitinous carapace, opisthosoma spines and a hard telson while the ventral section has soft tissues. Predators, like shorebirds, teleost, crabs, shrimp, crow (*Corvus splendens*) and wild pig (*Sus scrofa*) only consume the crab's soft tissue and would discard the exoskeleton (Debnath & Choudhury, 1988; Pati & Dash, 2016). Tribal (*e.g.* Nolia, Santhal, Oran and Munda) communities in India that live along estuaries would use *T. gigas* exoskeleton (carcasses) as decorative, produce jewellery, extract medicated oils (rheumatism, muscle aches and arthritis; Fig. 3), develop health tonic broths (aphrodisiac, spondylosis, bronchitis, wrist rheumatism, pneumonia and intestinal colic) or, the exoskeleton is used in rituals (supernatural and superstitious) (Chatterji & Abidi, 1994; Saha, 2010; Tripathy *et al.*, 2013). In 1995-1996, an adult *T. gigas* is valued at RS. 25 (USD 0.33) but after the nationwide seasonal fishing ban in 1998, enforcement was strengthened on vulnerable species and horseshoe crabs were listed in Schedule IV of the Wildlife (Protection) Act 1972 (Tripathy *et al.*, 2013; Behera *et al.*, 2015; Shinoj & Ramachandran, 2017).

Now, horseshoe crabs are considered as by-catch that damage fishing nets in northeast Odisha and West Bengal. Removal on land or discarding nets containing horseshoe crabs have made these crabs vulnerable to beach stranding from which, dead crabs are gathered by fish farm workers to prepare them as fish meal or compost (Khan, 2003; Tripathy *et al.*, 2013; Fig. 4). Recently, sedimentation from land reclamation, removal of mangrove forests and sand mining are threatening horseshoe crab spawning grounds (Tripathy *et al.*, 2013). Interventions, like rip-rap, Groyne and wave-breakers, were



(a)



(b)



(c)

Figure 4: a. Discarded (by-catch) juvenile horseshoe crab. b. Entanglement in fishing nets resulting in the mortality of horseshoe crab. c. Trapped in between stone on constructed jetty constructed near spawning ground

introduced but submerged horseshoe crab spawning grounds, exposed them to high energy currents and therefore, these affected areas are no longer suitable for horseshoe crab spawning (Mishra *et al.*, 2015). In addition, discarded fishing nets on the shore and ghost nets are stranding and suffocating horseshoe crabs (Pati

et al., 2017). Although capture fisheries may not threaten horseshoe crabs in India, human activities in coastal zones are challenging the survival of adults, hatching success of trilobite larvae and also larvae growing into adults which requires 7-11 years of endurance in their natal grounds.

Present Actions in Horseshoe Crab Conservation

Demand for live horseshoe crabs in Maharashtra (West) and Andhra Pradesh (southeast) offers income opportunity (RS. 50 ~ USD 0.67 per crab) to artisanal fishers in northeast India. However, these crabs are retrieved as by-catch and sold to local importers on a very small scale because these middle-men arrive without prior notice (Pati et al., 2020c). Though tribal markets in Odisha have horseshoe crab by-products for sale, these communities either use carcasses or dead crabs which have become stranded. The Wildlife Protection Act 1972 only penalizes possession of more than 3 live crabs (personal consumption). Although tribal communities are exempted from this clause, they only use horseshoe crab carcasses for their cultural practices. With fishing season bans (61 days) and fishing gear controls (cast net, bottom trawler net and gill net for shellfish) governed by the Fisheries Act (1897), regulations on net soaking and catch-per-unit gives horseshoe crabs an opportunity to reproduce and sustain their wild populations. Therefore, the only delimiting impact on horseshoe crabs is by-catch. Awareness spreading through questionnaires (2017-2018) have convinced 388 individuals to carefully remove horseshoe crabs from fishing nets and return them to the sea. A series of sea ranching projects in 2017-2019 that involved registered volunteers from the Association for Biodiversity Conservation and Research (ABC) who collected the eggs, incubate them in a closed hatchery setup and released the hatched (95 % hatching success) instar 1 larvae during lunar ebb tides have successfully recruited over 7,800 *T. gigas* larvae into Balaramgadi and Chandipur (Odisha). By far, horseshoe

crab conservation is in the implementing stage because it is challenged by area of coverage and communication. Also, volunteer training is time consuming and thus, the current manpower would require at least another five years to successfully spread and implement sea ranching outreach to the entire northeast coast of India.

References

- Annandale, N. (1909). The habitats of Indian king crabs. *Record of the Indian Museum*, 3, 294-295.
- Bannon, G. A., & Brown, G. G. (1980). Ultrastructural characteristics of the non-expanded and expanded extra-embryonic shell of the horseshoe crab, *Limulus polyphemus*L. *The Biological Bulletin*, 159(3), 582-591.
- Battelle, B. A., Calman, B. G., Andrews, A. W., Grieco, F. D., Mleziva, M. B., Callaway, J. C., & Stuart, A. E. (1991). Histamine: A putative afferent neurotransmitter in *Limulus* eyes. *Journal of Comparative Neurology*, 305(4), 527-542.
- Behera, S., Tripathy, B., Sivakumar, K., Choudhury, B. C., & Bhadury, P. (2015). Distribution and abundance of two sympatric species of horseshoe crabs along the Odisha Coast, India. In *Changing Global Perspectives on Horseshoe Crab Biology, Conservation and Management* (pp. 181-191). Cham: Springer.
- Bennett, D. C., Pezalla, P. D., & Herman, W. S. (1972). A sexual mosaic with internal embryogenesis in the dioecious horseshoe crab, *Limulus polyphemus*. *Bijdragen tot de Dierkunde*, 42(2), 192-192.
- Biswal, G. C., Andia, B. N., Pati, S., & Dash, B. P. (2016). Conservation of Indian horseshoe crab, *Tachypleus gigas* through captive rearing. *Frontiers in Life Sciences*, 178-181.
- Chatterji, A., & Pati, S. (2014). Allometric relationship in the adult population of the Malaysian horseshoe crab (*Tachypleus*

- tridentatus*; Leach). *International Journal of Research*, 1(11), 1378-1385.
- Chatterji, A., & Shaharom, F. (2009). Unusual spawning behaviour of the horseshoe crab (*Tachypleus gigas*, Muller) after the Tsunami along Orissa coast, India. *Pertanika Journal of Science and Technology*, 17(2), 263-268.
- Chatterji, A., Alam, H. A., Joshi, K. B., & Bhone, R. R. (2004a). Regeneration of gill lamellae of the Indian Horseshoe crab, *Tachypleus gigas* (Muller). *Current Science*, 87(11), 1511-1512.
- Chatterji, A., Kotnala, S., & Mathew, R. (2004b). Effect of salinity on larval growth of horseshoe crab, *Tachypleus gigas* (Muller). *Current Science*, 87(2), 248-250.
- Chatterji, A. (1999). New record of the sympatric distribution of two Asian species of the horseshoe crab. *Current Science*, 77(6), 746-747.
- Chatterji, A., Agular, Q., & Saldanha, C. (1996a). Energy source in the developing eggs of the Indian horseshoe crab, *Tachypleus gigas* (Muller). *Journal of Aquaculture in Tropic*, 11, 271-276.
- Chatterji, A., Parulekar, H. A., & Qasim, Z. (1996b). Nesting behaviour of the Indian horseshoe crab, *Tachypleus gigas* (Muller) (Xiphosura). In S. Z. Qasim, G. S. Roonwal (Eds.), *India's Exclusive Economic Zone* (pp. 142-152). India: Omega Scientific Publishers.
- Chatterji, A. (1994). The horseshoe crab-A living fossil. *A Project Swarajya Publication*, 157pp.
- Chatterji, A., & Abidi, S. A. H. (1994). The Indian horseshoe crab-A living fossil. *Journal of Indian Ocean Study*, 1(1), 42-48.
- Chatterji, A., Mishra, J. K., Vijayakumar, R., & Parulekar, H. A. (1994). Length-weight relationship of the Indian horseshoe crab *Tachypleus gigas* (Muller). *Indian Journal of Fisheries*, 41(2), 111-113.
- Chatterji, A., Vijayakumar, R. A., & Parulekar, H. A. (1992a). Spawning migration in of the horseshoe crab, *Tachypleus gigas* (Muller), in relation to the lunar cycle. *Asian Fisheries Science*, 5, 123-128.
- Chatterji, A., Vijayakumar, R. A., & Parulekar, H. A. (1992b). Seasonal variation in the volume of the hemolymph and body weight of *Tachypleus gigas* (Muller). *Pakistan Journals of Scientific and Industrial Research*, 35, 495-498.
- Chatterji, A., Mishra, J. K., & Parulekar, H. A. (1992c). Feeding behaviour and food selection in the horseshoe crab, *Tachypleus gigas* (Müller). *Hydrobiologia*, 246(1), 41-48.
- Chatterji, A., & Parulekar, H. A. (1992). Fecundity of the Indian horseshoe crab, *Carcinoscorpius rotundicauda* (Latreille). *Tropical Ecology*, 33(1), 97-102.
- Chatterji, A., Vijayakumar, R. A., & Parulekar, H. A. (1988). Growth and morphometric characteristics in the horseshoe crab, *Carcinoscorpius rotundicauda* (Latreille) from canning (West Bengal), India. *Pakistan Journals of Scientific and Industrial Research*, 31(5), 352-353.
- Chinnari, S., Pati, S., Dash, B. P., & Chatterji, A. (2015). A new record on inducement of differentiation of dendritic cells by the perivitelline fluid of the fertilized eggs of Indian horseshoe crab (*Tachypleus gigas*, Müller). *Biological forum-An International Journal*, 7(2), 678-681.
- Chiu, H. M., & Morton, B. (2003). Growth and allometry of two horseshoe crab species, *Tachypleus tridentatus* and *Carcinoscorpius rotundicauda* (Xiphosura), in Hong Kong. *Asian Marine Biology*, 18(2001), 129-141.
- Debnath, R., & Choudhury, A. (1992). Rare occurrence of tandem amplexus by the male Indian horseshoe crabs, *Carcinoscorpius rotundicauda*, with

- a comment on mating in imulidae (Merostomata: Xiphosura). *Advance in the Biosciences*, 15, 103-107.
- Debnath, R. (1991). Morphology and sexual dimorphism of Indian horseshoe crabs. *Geobios Newreport*, 10, 128-131.
- Debnath, R., & Choudhury, A. (1991). Comparative assessment of two breeding populations of crabs *Tachypleus gigas* (Miller) (Merostomata: Xiphosura). *Environment and Ecology*, 9, 163-166.
- Debnath, R., Nag, K. S., Choudhury, A., Dasgupta, R., & Sur, K. (1989). Feeding habit and digestive physiology of the Indian horseshoe crab, *Tachypleus gigas* (Muller). *Indian Journal of Physiological and Allied Science*, 43(2), 44-49.
- Debnath, R., & Choudhury, A. (1988). Population estimation of horseshoe crab *Tachypleus gigas* by capture-recapture method at Chandipur sea shore (Orissa), India. *Journal of Marine Biological Association of India*, 30, 8-12.
- Debnath, R. (1985). Studies on the distribution, abundance, breeding behaviour and feeding habit of Indian horseshoe crabs (Merostomata: Xiphosura) in the coastal water of Orissa and West Bengal, India. M. Phil. Dissertation, University of Calcutta, Calcutta (unpublished).
- Dumont, J. N., Anderson, E., & Chomyn, E. (1965). The anatomy of the peripheral nerve and its ensheathing artery in the horseshoe crab, Xiphosura (*Limulus polyphemus*). *Journal of Ultrastructure Research*, 13(1-2), 38-64.
- Fairuz-Fozi, N., Satyanarayana, B., Zauki, N. A. M., Muslim, A. M., Husain, M. L., Ibrahim, S., & Nelson, B. R. (2018). *Carcinoscorpius rotundicauda* (Latreille, 1802) population status and spawning behaviour at Pendas coast, Peninsular Malaysia. *Global Ecology and Conservation*, 15, e00422.
- Ghaskadbia, S., Patwardhan, V., Chakraborty, M., Agrawala, S., Verma, K. M., Chatterji, A., Lenka, N., & Parba, B. P. (2008). Enhancement of vertebrate cardiogenesis by a lectin from perivitelline fluid of horseshoe crab embryo. *Cellular and Molecular Life Science*, 65, 3312-3324.
- Graham, L. J., Murphy, B. R., & Hata, D. (2009). Using species composition data from a trawl survey to determine potential bycatch of the commercial trawl fishery for horseshoe crab *Limulus polyphemus* in the Middle Atlantic Bight. *North American Journal of Fisheries Management*, 29(2), 478-487.
- Heinbecker, P. (1932). The heart and median cardiac nerve of *Limulus polyphemus*. *American Journal of Physiology-Legacy Content*, 103(1), 104-120.
- Jawahir, A. R. N., Samsur, M., Shabdin, M. L., & Rahim, K. A. A. (2017). Morphometric allometry of horseshoe crab, *Tachypleus gigas* at west part of Sarawak waters, Borneo, East Malaysia. *Aquaculture, Aquarium, Conservation & Legislation*, 10(1), 18-24.
- John, B. A., Nelson, B. R., Sheikh, H. I., Cheung, S. G., Wardiatno, Y., Dash, B. P., Tsuchiya, K., Iwasaki, Y., & Pati, S. (2018). A review of fisheries and conservation status of Asian horseshoe crab. *Biodiversity and Conservation*, 27, 3573-3598.
- Joshi, B., Chatterji, A., & Bhonde, R. (2002). Long-term in vitro generation of Amoebocytes from the Indian horseshoe crab *Tachypleus gigas* (Muller). *In vitro cellular and Developmental Biology Animal*, 38(5), 255-257.
- Khan, R. A. (2003). Observation on some aspects of the biology of horseshoe crab, *Carcinoscorpius rotundicauda* (Latreille) on mud flats of Sundarban estuarine region. *Records of Zoological Survey of India*, 101(1&2), 1-23.
- Kumar, V., Roy, S., Sahoo, A. K., Behera, B. K., & Sharma, A. P. (2015). Horseshoe crab and its medicinal values. *International Journal of Current Microbiology and Applied Sciences*, 4(2), 956-964.

- Lazarus, S., Nammalwar, P., Pillai, V. N., Devadoss, P., & Mohanraj, G. (1990). Occurrence of king crab, *Tachypleus gigas* (Muller), off the Northeast coast of India. *Proceeding of the first Workshop on Scientific Results of FORV, Sagar Sampada, 1989*, 393-395.
- Levin, T. M. (2013). *Evidence for the existence of juvenile hormone in the horseshoe crab* (Doctoral dissertation, Worcester Polytechnic Institute), pp. 59-176.
- Mirshahi, M., Alam, H., Mirshahi, P., Soria, J., Therwath, A., & Chatterji, A. (2011). Role of Peri vitelline fluid of the fertilized eggs of Indian horseshoe crab promoting wound healing process. *Journal of Coastal Environment*, 2(2), 159-166.
- Mishra, J. K., Mishra, A., & Yasmin, Y. (2015). Status and threat perceptions of the Indian horseshoe crabs along the northeast coast of Bay of Bengal, India. In Carmichael, R. H., Botton, M. L., Shin, P. K. S., & Cheung, S. G. (Eds.), *Changing global perspectives on horseshoe crab biology, conservation and management* (pp. 397-406). Switzerland: Springer.
- Mishra, J. K. (2009a). Horseshoe crabs, their eco-biological status along the northeast coast of India and the necessity for ecological conservation. In Tanacredi, J. T., Botton, M. L., Smith, D. (Eds.), *Biology and conservation of horseshoe crabs* (pp. 89-96). Boston, MA: Springer.
- Mishra, J. K. (2009b). Larval culture of *Tachypleus gigas* and its molting behaviour under laboratory conditions. In Tanacredi, J. T., Botton, M. L., Smith, D. (Eds.), *Biology and conservation of horseshoe crabs* (pp. 513-519). Springer.
- Mishra, J. K., Chatterji, A., & Parulekar, H. A. (1993). A freak twin larva of the Indian horseshoe crab *Tachypleus gigas* (Muller). *Journal of Bombay Natural History Society*, 90(1), 116-117.
- Modayil, M. J., Sathiadhas, R., & Gopakumar, G. (2006). Marine farming: Country Analysis-India. In *FAO/NACA Regional Workshop on the Future of Mariculture: A Regional Approach for Responsible Development in the Asia-Pacific Region* (pp. 7-11). Guangzhou, China.
- Mohamad, F., Manca, A., Ahmad, A., & Fawwaz, M. (2016). Width-weight and length-weight relationships of the tri-spine horseshoe crab, *Tachypleus tridentatus* (leach 1819) from two populations in Sabah, Malaysia: implications for population management. *Journal of Sustainability Science and Management*, 11(1), 1-13.
- Mondal, I., & Bandyopadhyay, J. (2014). Coastal zone mapping through geospatial technology for resource management of Indian Sundarban, West Bengal, India. *International Journal of Remote Sensing Applications*, 4(2), 103-112.
- Nair, S. R., Parulekar, H. A., & Desai, B. N. (1990). Research in the assessment of capture and culture fisheries along the Indian coast. *Central Marine Fisheries Research Institute Bulletin*, 44(2), 297-305.
- Nelson, B. R., John, A., Zhong, J. H., & Jalal, K. C. A. (2020). Membrane retention potential of *Tachypleus gigas* during early embryogenesis. *Desalination and Water Treatment*, 187, 17-23.
- Nelson, B. R., Zhong, J. M. H., Zauki, N. A. M., Satyanarayana, B., & Chowdhury, A. J. K. (2019). Effects of shore sedimentation to *Tachypleus gigas* (Müller, 1785) spawning activity from Malaysian waters. *Journal of Sustainability Science and Management*, 14(1), 41-60.
- Nelson, B. R., Satyanarayana, B., Moh, J. H. Z., Ikhwanuddin, M., Chatterji, A., & Shaharom, F. (2016a). The final spawning ground of *Tachypleus gigas* (Müller, 1785) on the east Peninsular Malaysia is at risk: A call for action. *PeerJ*, 4, e2232.
- Nelson, B. R., Satyanarayana, B., Moh, J. H. Z., & Shaharom, F. (2016b). Does human infringement at the spawning grounds challenge horseshoe crab eggs and their

- embryogenesis. *Journal of Sustainability Science and Management*, 1,1-10.
- Nelson, B. R., Satyanarayana, B., Zhong, J. M. H., Shaharom, F., Sukumaran, M., & Chatterji, A. (2015). Episodic human activities and seasonal impacts on the *Tachypleus gigas* (Müller, 1785) population at Tanjung Selangor in Peninsular Malaysia. *Estuarine, Coastal and Shelf Science*, 164, 313-323.
- Nong, W., Qu, Z., Li, Y., Owen, T. B., Wong, A. Y., Yip, H. Y., ... & Cao, J. (2020). Horseshoe crab genomes reveal the evolutionary fates of genes and microRNAs after three rounds (3R) of whole genome duplication. *bioRxiv*.
- Oppenheim, J. D., Nachbar, M. S., Salton, M. R., & Aull, F. (1974). Purification of a hemagglutinin from *Limulus polyphemus* by affinity chromatography. *Biochemical and Biophysical Research Communications*, 58(4), 1127-1134.
- Panda, S., & Naik, P. K. (2017). Morphometric study of horseshoe crab (*Carcinoscorpius rotundicauda*) in Odisha. *Journal of Surgery*, 3(2), 48-53.
- Pati, S., Shahimi, S., Nandi, D., Sarkar, T., Acharya, S. N., Sheikh, H. I. ... Atan, E. H. (2021). Predicting *Tachypleus gigas* spawning distribution with climate change in Northeast Coast of India. *Journal of Ecological Engineering*.
- Pati, S., Jena, P., Shahimi, S., Nelson, B. R., Acharya, D., Dash, B. P., & Chatterji, A. (2020a). Characterization dataset for pre-and post-irradiated shrimp waste chitosan. *Data in Brief*, 106081.
- Pati, S., Chatterji, A., Dash, B. P., Nelson, B. R., Sarkar, T., Shahimi, S., Edinur, H. A., Binti Abd Manan, T.S., Jena, P., Mohanta, Y. K., & Acharya, D. (2020b). Structural characterization and antioxidant potential of Chitosan by γ -Irradiation from the carapace of horseshoe crab. *Polymers*, 12(10), 2361.
- Pati, S., Shahimi, S., Atan Edinur, H., Acharya, D., Dash, B. P., & Nelson, B. R. (2020c). Extraction of people's perception towards horseshoe crab existence in Northeast India. *Frontiers in Marine Science*, 7, 924.
- Pati, S., Chatterji, A., & Dash, B. P. (2018). Chitosan from the carapace of Indian horseshoe crab (*Tachypleus gigas*, Müller): Isolation and its characterization. *Advances in BioResearch*, 9(4), 52-64.
- Pati, S., Tudu, S., Rajesh, A., Biswal, G. C., Chatterji, A., Dash, B. P., & Samantaray, R. (2017). Man-made activities affecting the breeding ground of horseshoe crab, *Tachypleus gigas* (Müller, 1795) along Balasore coast: Call for immediate conservation. *E-planet*, 15(2), 145-154.
- Pati, S., & Dash, B. P. (2016). Horseshoe crab (*Tachypleus gigas*) as prey of domestic pig (*Sus domesticus*) in Khandia estuary, Balasore, Odisha, India. *Zool Print*, 31(5), 14-15.
- Pati, S., Biswal, G. C., & Dash, B. P. (2015). Availability of *Tachypleus gigas* (Müller) along the river estuaries of Balasore district, Odisha, India. *International Journal of Fisheries and Aquatic Studies*, 2, 334-336.
- Patil, S. J., & Anil, C. A. (2000). Epibiotic community of the horseshoe crab *Tachypleus gigas*. *Marine Biology*, 136, 699-713.
- Putri, W. A., Purwiyanto, A. I., Agustriani, F., & Mustopa, A. Z. (2019). The morphometric variability of the mangrove horseshoe crab (*Carcinoscorpius rotundicauda*) from Banyuasin estuarine of South Sumatra, Indonesia. *Ecologica Montenegrina*, 24, 38-46.
- Rao, K.V. R., & Rao, K. V. S. (1972). Studies on Indian king crab (Arachnida: Xiphosura). *Proceeding of Indian National Science Academy*, 38B(3/4), 206-211.
- Razak, M. R., & Kassim, Z. (2018). Comparison of horseshoe crabs (*Tachypleus gigas*) morphometry between different populations using allometric analysis. *Aquaculture*,

- Aquarium, Conservation & Legislation*, 11(1), 143-157.
- Roonwal, M. L. (1944). Some observations on the breeding biology, and on the swelling, weight, water content and embryonic movements in the developing, eggs, of the Moluccasking crab, *Tachypleus gigas* (Muller) (Arthropoda: Xiphosura). *Proceeding of the National Academic of science*, 1 (Sec-B 20), 115-129.
- Saha, D. (2010). Ecological importance of horseshoe crabs. *The Ecological Research and Development*, 2(1), 158-162.
- Sahu, A. C., & Dey, L. (2013). Spawning density and morphometric characteristics of the Horseshoe crab *Tachypleus gigas*(Muller) on the Balasore coast of Bay of Bengal. *India Science Vision*, 13(2), 76-84.
- Seawall, S. B. R. (1912). Xiphosura. *Record of the Indian Museum*, 7, 87-88.
- Sekiguchi, K., Nakamura, K., Sen, T. K., & Sugita, H. (1976). Morphological variation and distribution of a horseshoe crab, *Tachypleus gigas*, from the Bay of Bengal and Gulf of Siam. *Proceeding of The Japanese Society of Systematic Zoology*, 12, 13-20.
- Shinoj, P., & Ramachandran, C. (2017). Taming the fishing blues reforming the marine fishery regulatory regime in India. *Economic and Political Weekly*, 52(45), 73-81.
- Shuster Jr., C. N. (1948). *On the gross anatomy and histology of the alimentary tract in early developmental stages of Xiphosura (= Limulus) polyphemus Linn.* MS thesis (pp. 37-93), Rutgers University, New Jersey.
- Smith, M. D., & Brockmann, H. J. (2014). The evolution and maintenance of sexual size dimorphism in horseshoe crabs: An evaluation of six functional hypotheses. *Animal Behaviour*, 96, 127-139.
- Srijaya, T. C., Pradeep, P. J., Mithun, S., Hassan, A., Shaharom, F., & Chatterji, A. (2010). A new record on the morphometric variations in the populations of horseshoe crab (*Carcinoscorpius rotundicauda* Latreille) obtained from two different ecological habitats of Peninsular Malaysia. *Our Nature*, 8(1), 204-211.
- Srimal, S., Miyata, T., Kawabata, S., Miyata, T., & Iwanaga, S. (1985). The complete amino acid sequence of coagulogen isolated from southeast Asian horseshoe crab, *Carcinoscorpius rotundicauda*. *Journal of Biochemistry*, 98, 305-318.
- Syuhaida, N., Rozihan, M., Akbar, J., Akmal, M., & Joni, H. (2019). Allometry relationship of mangrove horseshoe crab, *Carcinoscorpius rotundicauda* from the West Coast of Peninsular Malaysia. *International Journal of Fisheries and Aquatic Studies*, 7(2), 223-228.
- Techera, E. (2020). Indian Ocean fisheries regulation: exploring participatory approaches to support small-scale fisheries in six States. *Journal of the Indian Ocean Region*, 16(1), 27-46.
- Tripathy, B., Sivakumar, K., Johan, S., Behera, S., & Choudhury, C. B. (2013). Assessment of the population status and threats to the horseshoe crabs along the northern east coast of India, p. 137-146. In *Ecology and conservation of Tropical Marine Faunal communities*, Springer, 481p.
- Vijayakumar, R., Das, S., Chatterji, A., & Parulekar, A. (2000). Morphometric characteristics in the horseshoe crab *Tachypleus gigas* (Arthropoda: Merostomata). *Indian Journal of Marine Sciences*, 29(4), 333-335.
- Walls, E. A., Berkson, J., & Smith, S. A. (2002). The horseshoe crab, *Limulus polyphemus*: 200 million years of existence, 100 years of study. *Reviews in Fisheries Science*, 10(1), 39-73.
- Weaver, L. T., Freiberg, E., Israel, E. I., & Walker, W. A. (1988). Epidermal growth factor in human amniotic fluid. *Gastroenterology*, 95(5), 1436.

- Yamamichi, Y., Sugita, H., & Sekiguchi, K. (1983). Morphological characterization of first instar larvae of Asian horseshoe crabs and their hybrids* (horseshoe crabs/ interspecific hybrids/larvae). *Development, Growth & Differentiation*, 25(3), 271-280.
- Yamamichi, Y., & Sekiguchi, K. (1974). Embryo and organ cultures of the horseshoe crab, *Tachypleus tridentatus*. *Development, Growth & Differentiation*, 16(4), 295-304.
- Yennawar, P. (2015). Status of horseshoe crab at Digha, Northern east coast of India. P. 89-93. In *Marine Faunal Diversity in India*, (519p). Elsevier.
- Zauki, N. A. M., Satyanarayana, B., Fairuz-Fozi, N., Nelson, B. R., Martin, M. B., Akbar-John, B., & Chowdhury, A. J. K. (2019a). Citizen science frontiers horseshoe crab population regain at their spawning beach in East Peninsular Malaysia. *Journal of Environmental Management*, 232, 1012-1020.
- Zauki, N. A. M., Satyanarayana, B., Fairuz-Fozi, N., Nelson, B. R., Martin, M. B., Akbar-John, B., & Chowdhury, A. J. K. (2019b). Horseshoe crab bio-ecological data from Balok, east coast peninsular Malaysia. *Data in Brief*, 22, 458-463.